

Procidic[®]

Broad Spectrum Bactericide and Fungicide Compound

Fruits and Vegetables Crops

Your solution for fungal & bacterial borne diseases.

Procidic is a bactericide and fungicide formulated to protect crops during all stages of growth. Procidic's advanced formula works to eliminate pathogenic bacteria and fungi on contact and is rapidly absorbed into the plant.

Working systemically, Procidic moves toward the apex of the plant to inhibit the future growth of bacteria and fungi. Procidic may be applied in regular foliar spray programs or in tank mixes. Always apply according to the directions on the label.

Procidic is the IDEAL solution for rotation practices as part of your fungicide cross resistance management program.

Prevents and Controls

- Damping Off
- Downy Mildew
- Powdery Mildew
- Alternaria sp
- Botrytis sp
- Colletotrichum sp
- Angular Leaf Spot
- Pseudomonas sp

- Bactericide and fungicide dual action mode
- Organically certifiable
- Works systemically
- No re-entry waiting period required
- Works on contact & through residual action
- No days to harvest interval



See back for more details or visit us at www.procidic.com

ACTIVE INGREDIENTS	
Citric acid.....	3.5%
OTHER INGREDIENTS *.....96.5%	
TOTAL	100%

*Vitamin C, ethyl lactate, glycerin, sodium chloride, water

Greenspire Global, Inc. represents that this product is a minimum risk pest control product and qualifies for exemption from EPA registration under FFIFRA25 (b).

Procidic®

Broad Spectrum Bactericide and Fungicide Compound

Recommended Usage Rates & Instructions

1. Procidic is best used as a preventative.
2. Apply Procidic in regularly scheduled spray program.
3. Procidic requires quality water free of contaminants and organic matter, a pH of 5.5-6.0 and a low salt concentration.
4. No re-entry waiting period is required.
5. Use higher dosage of Procidic during periods of high humidity.
6. Refrain from unnecessary contact with crops after treatment.
7. Do not mix with dicofol, methomyl and cyhexatin.
8. Do not mix with salt-based or high-alkali reaction products.
9. Procidic is absorbed systemically through the stem, leaves and root system. When disease pressure is present Procidic applications must reach live, green leaves.



Use Rates



CUCURBITS: Watermelon, melon, pumpkin, zucchini, squash

Disease Downy Mildew (*Pseudoperonospora cubensis*),
Powdery Mildew (*Sphaerotheca fulginea*)

Dose 3.0 fl oz/acre

Use Rate After transplanting, spray day 12 and days 20-22.

Dose 4.5 fl oz/acre

Use Rate After transplanting, spray during days 28-30.

Dose 6 fl oz/A

Use Rate After transplanting, spray during days 32-34 and days 40-42.

Disease Damping Off (*Fusarium sp*, *Pythium sp*)

Dose 7-11 fl oz/acre

Use Rate After transplanting, spray during days 32-34 and days 40-42.

CUCUMBER

Disease Bacterial Wilt of Cucumber

Dose 3.5 fl oz/acre

Use Rate After transplanting, spray during day 12 and during days 20-22.

► For more information go to www.procidic.com or call 877.978.6558